



## BAD (Phospho-Ser136) Antibody

#11068

**Catalog Number:** 11068-1, 11068-2

**Amount:** 50  $\mu$ g/50  $\mu$ l, 100  $\mu$ g/100  $\mu$ l

**Swiss-Prot No. :** Q61337

**Form of Antibody:** Rabbit IgG in phosphate buffered saline (without  $Mg^{2+}$  and  $Ca^{2+}$ ), pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.

**Storage/Stability:** Store at  $-20^{\circ}C$ /1 year

**Immunogen:** The antiserum was produced against synthesized phosphopeptide derived from mouse BAD around the phosphorylation site of serine136 (S-R-S<sub>P</sub>-A-P).

**Purification:** The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific phosphopeptide. The antibody against non-phosphopeptide was removed by chromatography using non-phosphopeptide corresponding to the phosphorylation site.


**Specificity/Sensitivity:** BAD (phospho-Ser136) antibody detects endogenous levels of BAD only when phosphorylated at Serine 136.

**Reactivity:** Human, Mouse, Rat


### Applications:

Predicted MW: 23kd

WB: 1:500~1:1000 IHC: 1:50~1:100



Western blot analysis of extracts from HT29 cells untreated or treated with UV using BAD(Phospho-Ser136) Antibody #11068.



P-Peptide -

+

Immunohistochemical analysis of paraffin-embedded human breast carcinoma tissue using BAD (phospho-Ser136) antibody (#11068).

### Background :

The protein encoded by BAD gene is a member of the BCL-2 family. BCL-2 family members are known to be regulators of programmed cell death. This protein positively regulates cell apoptosis by forming heterodimers with BCL-xL and BCL-2, and reversing their death repressor activity. Proapoptotic activity of this protein is regulated through its phosphorylation. Protein kinases AKT and MAP kinase, as well as protein phosphatase calcineurin were found to be involved in the regulation of this protein. Alternative splicing of this gene results in two transcript variants which encode the same isoform.

**References:** N Matsuda, Y Takano, S Kageyama, et al. (2007) Silencing of caspase-8 and caspase-3 by

RNA interference prevents vascular endothelial cell injury in mice with endotoxic shock. Cardiovascular Research, 76:132-140.

This article references the use of the **#11068** in the following applications: **Western blotting**