
Order : order@swbio.com

#11069



Catalog Number: 11069-1, 11069-2

Amount: 50 μ g/50 μ 1, 100 μ g/100 μ 1

Swiss-Prot No. : Q61337

Form of Antibody: Rabbit IgG in phosphate buffered saline (without Mg2+ and Ca2+), pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.

Storage/Stability: Store at −20°C/1 year

Immunogen: The antiserum was produced against synthesized phosphopeptide derived from mouse BAD around the phosphorylation site of serine 155 (R-M-Sp-D-E).


Purification: The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific phosphopeptide. The antibody against non-phosphopeptide was removed by chromatography using non-phosphopeptide corresponding to the phosphorylation site.

Specificity/Sensitivity: BAD (phospho-Ser155) antibody detects endogenous levels of BAD only when phosphorylated at serine 155.

Reactivity: Human, Mouse, Rat

Applications:

Predicted MW: 23kd WB:1:500~1:1000 IHC:1:50~1:100




Forskolin + +

P-Peptide - +

Western blot analysis of extracts from 293 cells using

BAD (phospho-Ser155) antibody (#11069).



P-Peptide - + Immunohistochemical analysis of paraffin-embedded human breast carcinoma tissue using BAD (phospho-Ser155) antibody (#11069).

Background :

The protein encoded by BAD gene is a member of the BCL-2 family. BCL-2 family members are known to be regulators of programmed cell death. This protein positively regulates cell apoptosis by forming heterodimers with BCL-xL and BCL-2, and reversing their death repressor activity. Proapoptotic activity of this protein is regulated through its phosphorylation. Protein kinases AKT and MAP kinase, as well as protein phosphatase calcineurin were found to be involved in the regulation of this protein. Alternative splicing of this gene results in two transcript variants which encode the same isoform.

References: Moon EY, et al. (2003). Blood. 101(10): 4122-4130.