

## Histone H3 (Di Methyl Lys27) (3B12)



## Mouse monoclonal Antibody

Catalog Number: 55206 Amount: 100µg/100µl

Swiss-Prot No.: P68431/Q71DI3/P84243

GENE ID:8350/8351/8352/8353/8354/8355/8356/8357/8358/8968

Form of Antibody: Purified mouse monoclonal in buffer containing 0.1M Tris-Glycine (pH 7.4, 150 mM

NaCl) with 0.2% sodium azide, 50%,glycerol **Storage/Stability:** Store at -20°C/1 year **Purification:** affinity-chromatography

Specificity/Sensitivity: This antibody detects endogenous levels of Histone H3 (Di Methyl Lys27) and does


not corss-react with related proteins

**Alternative Names:**HIST1H3A; H3FA; HIST1H3B; H3FL; HIST1H3C; H3FC; HIST1H3D; H3FB; HIST1H3E; H3FD; HIST1H3F; H3FI; HIST1H3G; H3FH; HIST1H3H; H3FK; HIST1H3I; H3FF; HIST1H3J; H3FJ; Histone H3.1; Histone H3/a; Histone H3/b; Histone H3/c; Histone H3/d; Histone H3/f; Histone H3/

Reactivity: Human, Mouse, Rat

## **Applications:**

Predicted MW: 15kd WB:1:1000-3000 IP:1:100-200



Western blot analysis of Hela using Histone H3 (Di Methyl Lys27) Mouse mAb

**Background:**Core component of nucleosome. Nucleosomes wrap and compact DNA into chromatin, limiting DNA accessibility to the cellular machineries which require DNA as a template. Histones thereby play a central role in transcription regulation, DNA repair, DNA replication and chromosomal stability. DNA accessibility is regulated via a complex set of post-translational modifications of histones, also called histone code, and nucleosome remodeling